

Algebra $\mathfrak{gl}(\lambda)$ Inside the Algebra of Differential Operators on the Real Line

H GARGOUBI

I.P.E.I.M., route de Kairouan, 5019 Monastir, Tunisia

E-mail: hichem.gargoubi@ipeim.rnu.tn

Received November 22, 2001; Revised February 26, 2002; Accepted March 12, 2002

Abstract

The Lie algebra $\mathfrak{gl}(\lambda)$ with $\lambda \in \mathbb{C}$, introduced by B L Feigin, can be embedded into the Lie algebra of differential operators on the real line (see [7]). We give an explicit formula of the embedding of $\mathfrak{gl}(\lambda)$ into the algebra $\mathcal{D}_\lambda$ of differential operators on the space of tensor densities of degree λ on $\mathbb{R}$. Our main tool is the notion of projectively equivariant symbol of a differential operator.

1 Introduction

The Lie algebra $\mathfrak{gl}(\lambda)$ ($\lambda \in \mathbb{C}$) was introduced by B L Feigin in [7] for calculation the cohomology of the Lie algebra of differential operators on the real line. The algebra $\mathfrak{gl}(\lambda)$ is defined as the quotient of the universal enveloping algebra $U(\mathfrak{sl}_2)$ of $\mathfrak{sl}_2$ with respect to the ideal generated by the element $\Delta - \lambda(\lambda - 1)$, where Δ is the Casimir element of $U(\mathfrak{sl}_2)$. $\mathfrak{gl}(\lambda)$ is turned into a Lie algebra by the standard method of setting $[a, b] = ab - ba$.

According to Feigin, $\mathfrak{gl}(\lambda)$ can be considered as an analogue of $\mathfrak{gl}(n)$ for $n = \lambda \in \mathbb{N}$; it is also called the algebra of matrices of complex size, see also [13, 16, 17, 12].

We consider the space $\mathcal{D}_\lambda$ of all linear differential operators acting on tensor densities of degree λ on $\mathbb{R}$. One of the main results of [7] is the construction of an embedding $\mathfrak{gl}(\lambda) \rightarrow \mathcal{D}_\lambda$.

The purpose of this paper is to give an explicit formula of this embedding. We also show that this embedding realizes the isomorphism of Lie algebras $\mathfrak{gl}(\lambda) \cong \mathcal{D}_\lambda^{\text{pol}}$ constructed in [1, 2], where $\mathcal{D}_\lambda^{\text{pol}} \subset \mathcal{D}_\lambda$ is the subalgebra of differential operators with polynomial coefficients.

The main idea of this paper is to use the *projectively equivariant symbol* of a differential operator, that is an $\mathfrak{sl}_2$ -equivariant way to associate a polynomial function on $T^*\mathbb{R}$ to a differential operator. The notion of projectively equivariant symbol was defined in [4, 15] and used in [8, 9, 10] for study of modules of differential operators.

2 Basic definitions

2.1 The Lie algebra $\mathfrak{gl}(\lambda)$. Let $\text{Vect}(\mathbb{R})$ be the Lie algebra of smooth vector fields on $\mathbb{R}$ with complex coefficients: $X = X(x)\partial$, where $X(x)$ is a smooth complex function of one real variable; $X(x) \in C^\infty(\mathbb{R}, \mathbb{C})$, and where $\partial = \frac{d}{dx}$. Consider the Lie algebra $\mathfrak{sl}_2 \subset \text{Vect}(\mathbb{R})$ generated by the vector fields

$$\{\partial, x\partial, x^2\partial\}. \quad (2.1)$$

Denote $e_i := x^i\partial$, $i = 0, 1, 2$, the Casimir element

$$\Delta := e_1^2 - \frac{1}{2}(e_0e_2 + e_2e_0)$$

generates the center of $U(\mathfrak{sl}_2)$. The quotient

$$\mathfrak{gl}(\lambda) := U(\mathfrak{sl}_2)/(\Delta - \lambda(\lambda - 1)), \quad \lambda \in \mathbb{C}$$

is naturally a Lie algebra containing $\mathfrak{sl}_2$.

2.2 Modules of differential operators on $\mathbb{R}$. Denote $\mathcal{D}$ the Lie algebra of linear differential operators on $\mathbb{R}$ with complex coefficients:

$$A = a_n(x)\partial^n + a_{n-1}(x)\partial^{n-1} + \cdots + a_0(x), \quad (2.2)$$

with $a_i(x) \in C^\infty(\mathbb{R}, \mathbb{C})$.

For $\lambda \in \mathbb{C}$, $\text{Vect}(\mathbb{R})$ is embedded into the Lie algebra $\mathcal{D}$ by:

$$X \mapsto L_X^\lambda := X(x)\partial + \lambda X'(x). \quad (2.3)$$

Denote $\mathcal{D}_\lambda$ the $\text{Vect}(\mathbb{R})$ -module structure with respect to the adjoint action of $\text{Vect}(\mathbb{R})$ on $\mathcal{D}$. The module $\mathcal{D}_\lambda$ has a natural filtration: $\mathcal{D}_\lambda^0 \subset \mathcal{D}_\lambda^1 \subset \cdots \subset \mathcal{D}_\lambda^n \subset \cdots$, where $\mathcal{D}_\lambda^n$ is the module of n -th order differential operators (2.2).

Geometrically speaking, differential operators are acting on tensor densities, namely: $A : \mathcal{F}_\lambda \rightarrow \mathcal{F}_\lambda$, where $\mathcal{F}_\lambda$ is the space of tensor densities of degree λ on $\mathbb{R}$ (i.e., of sections of the line bundle $(T^*\mathbb{R})^{\otimes \lambda}$, $\lambda \in \mathbb{C}$), that is: $\phi = \phi(x)(dx)^\lambda$, where $\phi(x) \in C^\infty(\mathbb{R}, \mathbb{C})$.

It is evident that $\mathcal{F}_\lambda \cong C^\infty(\mathbb{R}, \mathbb{C})$ as linear spaces (but not as modules) for any λ . We use this identification throughout this paper. The Lie algebra structures of differential operators acting on the space of tensor densities and on the space of functions are also identified (see [8]).

The $\text{Vect}(\mathbb{R})$ -modules $\mathcal{D}_\lambda$ were considered by classics (see [3, 18]) and, recently, studied in a series of papers [5, 9, 8, 10, 14].

2.3 Principal symbol. Let $\text{Pol}(T^*\mathbb{R})$ be the space of functions on $T^*\mathbb{R}$ polynomial in the fibers. This space is usually considered as the space of symbols associated to the space of differential operators on $\mathbb{R}$.

Recall that the *principal symbol* of a differential operator is the linear map $\sigma : \mathcal{D} \rightarrow \text{Pol}(T^*\mathbb{R})$ defined by:

$$\sigma(A) = a_n(x)\xi^n,$$

where A is a differential operator (2.2) and ξ is the coordinate on the fiber.

One can also speak about the principal symbol of an element of $U(\mathfrak{sl}_2)$. Indeed, $U(\mathfrak{sl}_2)$ is canonically identified with the symmetric algebra $S(\mathfrak{sl}_2)$ as $\mathfrak{sl}_2$ -modules (see, e.g., [6, p.82]). Using the realization (2.1), the algebra $S(\mathfrak{sl}_2)$ can be projected to $\text{Pol}(T^*\mathbb{R})$. Therefore, one can define in a natural way the principal symbol on $S(\mathfrak{sl}_2)$.

Our goal is to construct an $\mathfrak{sl}_2$ -equivariant linear map $T_\lambda : U(\mathfrak{sl}_2) \rightarrow \mathcal{D}_\lambda$ which preserves the principal symbol, i.e., such that the following diagram commutes:

$$\begin{array}{ccc} U(\mathfrak{sl}_2) & \xrightarrow{T_\lambda} & \mathcal{D}_\lambda \\ \sigma \downarrow & & \downarrow \sigma \\ \text{Pol}(T^*\mathbb{R}) & \xrightarrow{id} & \text{Pol}(T^*\mathbb{R}) \end{array}$$

2.4 Projectively equivariant symbol. Viewed as a $\text{Vect}(\mathbb{R})$ -module, the space of symbols corresponding to $\mathcal{D}_\lambda$ has the form:

$$\text{Pol}(T^*\mathbb{R}) \cong \mathcal{F}_0 \oplus \mathcal{F}_1 \oplus \cdots \oplus \mathcal{F}_n \oplus \cdots. \quad (2.4)$$

The space of polynomials of degree $\leq n$ is a submodule of $\text{Pol}(T^*\mathbb{R})$ which we denote $\text{Pol}_n(T^*\mathbb{R})$.

The following result of [8] allows one to identify, for arbitrary λ , $\mathcal{D}_\lambda^n$ with $\text{Pol}_n(T^*\mathbb{R})$ as $\mathfrak{sl}_2$ -modules:

(i) There exists a unique $\mathfrak{sl}(2, \mathbb{R})$ -isomorphism $\sigma_\lambda : \mathcal{D}_\lambda^n \rightarrow \text{Pol}_n(T^*\mathbb{R})$ preserving the principal symbol.

(ii) σ_λ associates to each differential operator A the polynomial $\sigma_\lambda(A) = \sum_{p=0}^n \bar{a}_p(x) \xi^p$, defined by:

$$\bar{a}_p(x) = \sum_{j=p}^n \alpha_p^j a_j^{(j-p)}, \quad (2.5)$$

where the constants α_p^j are given by:

$$\alpha_p^j = \frac{\binom{j}{p} \binom{2\lambda-p}{j-p}}{\binom{j+p+1}{2p+1}}$$

(the binomial coefficient $\binom{\lambda}{j} = \lambda(\lambda-1)\cdots(\lambda-j+1)/j!$ is a polynomial in λ).

The isomorphism σ_λ is called the *projectively equivariant symbol map*. Its explicit formula was first found in [4, 15] in the general case of pseudo-differential operators on a one-dimensional manifold (see also [15] for the multi-dimensional case).

3 Main result

In this section, we give the main result of this paper. We adopt the following notations:

$$[L_{X_1}^\lambda L_{X_2}^\lambda \cdots L_{X_n}^\lambda]_+ := \sum_{\tau \in S_n} L_{X_{\tau(1)}}^\lambda \circ L_{X_{\tau(2)}}^\lambda \circ \cdots \circ L_{X_{\tau(n)}}^\lambda$$

for a symmetric n -linear map from $\text{Vect}(\mathbb{R})$ to $\mathcal{D}$ and

$$(X_1 X_2 \cdots X_n)_+ := \sum_{\tau \in S_n} X_{\tau(1)} X_{\tau(2)} \cdots X_{\tau(n)}$$

for a symmetric n -linear map from $\mathfrak{sl}_2$ to $U(\mathfrak{sl}_2)$, where S_n is the group of permutations of n elements and $X_i \in \mathfrak{sl}_2$.

Theorem 1. (i) For arbitrary $\lambda \in \mathbb{C}$, there exists a unique $\mathfrak{sl}_2$ -equivariant linear map preserving the principal symbol:

$$T_\lambda : U(\mathfrak{sl}_2) \rightarrow \mathcal{D}_\lambda$$

defined by

$$T_\lambda((X_1 X_2 \cdots X_n)_+) = [L_{X_1}^\lambda L_{X_2}^\lambda \cdots L_{X_n}^\lambda]_+, \quad (3.1)$$

where $X_i \in \{e_0, e_1, e_2\}$, $L_{X_i}^\lambda$ given by (2.3) and $n = 1, 2, \dots$.

(ii) The operator T_λ is given in term of the $\mathfrak{sl}_2$ -equivariant symbol (2.5) by:

$$\sigma_\lambda([L_{X_1}^\lambda L_{X_2}^\lambda \cdots L_{X_n}^\lambda]_+) = \sum_{\substack{0 \leq k \leq n \\ k \text{ even}}} P_k^n(\lambda) \mathcal{A}_k(X_1, \dots, X_n) \xi^{n-k}, \quad (3.2)$$

where

$$\begin{aligned} \mathcal{A}_k(X_1, \dots, X_n) \\ = \sum_{2p+m=k} \binom{k/2}{p} (-2)^p (X_1'' \cdots X_p'' X_{p+1}' \cdots X_{p+m}' X_{p+m+1} \cdots X_n)_+ \end{aligned} \quad (3.3)$$

and

$$P_k^n(\lambda) = \sum_{p=0}^n \sum_{l=n-k}^n (l-n+k)! \frac{\binom{l}{n-k}^2 \binom{2\lambda-n+k}{l-n+k}}{\binom{n-k+l+1}{2n-2k+1}} \binom{n}{p} \left\{ \begin{matrix} p \\ l \end{matrix} \right\} \lambda^{n-p}, \quad (3.4)$$

where $\left\{ \begin{matrix} p \\ l \end{matrix} \right\}$ is the Stirling number of the second kind¹.

It is worth noticing that the linear map T_λ does not depend on the choice of the PBW-base in $U(\mathfrak{sl}_2)$.

4 Proof of Theorem 1

By construction, the linear map T_λ is $\mathfrak{sl}_2$ -equivariant.

4.1 $\mathfrak{sl}_2$ -invariant symmetric differential operators. To prove part (ii) of Theorem 1 one needs the following

¹We refer to [11] as a nice elementary introduction to the combinatorics of the Stirling numbers.

Proposition 1. *For arbitrary $\mu \in \mathbb{C}$ and $n = 1, 2, \dots$, there exists at most one, up to proportionality, $\mathfrak{sl}_2$ -equivariant symmetric operator $\otimes^n \mathfrak{sl}_2 \rightarrow \mathcal{F}_\mu$ which is differential with respect to the vector fields $X_i \in \mathfrak{sl}_2$. This operator exists if and only if $\mu = k - n$, where k is an even positive integer. It is denoted: $\mathcal{A}_k : \otimes^n \mathfrak{sl}_2 \rightarrow \mathcal{F}_{k-n}$, and defined by the expression (3.3).*

Proof. Each k -th order differential operator $\mathcal{A} : \otimes^n \mathfrak{sl}_2 \rightarrow \mathcal{F}_\mu$ is of the form:

$$\mathcal{A}(X_1, \dots, X_n) = \sum_{2p+m=k} \beta_p(x) (X_1'' \cdots X_p'' X_{p+1}' \cdots X_{p+m}' X_{p+m+1} \cdots X_n)_+,$$

where $\beta_p(x)$ are some functions.

The condition of $\mathfrak{sl}_2$ -equivariance for $\mathcal{A}$ reads as follows:

$$X[\mathcal{A}(X_1, \dots, X_n)]' + \mu X' \mathcal{A}(X_1, \dots, X_n) = \sum_{i=1}^n \mathcal{A}(X_1, \dots, L_X^{-1}(X_i), \dots, X_n),$$

where $X \in \mathfrak{sl}_2$.

Substitute $X = \partial$ to check that the coefficients $\beta_p(x)$ do not depend on x . Substitute $X = x\partial$ to obtain the condition $\mu = k - n$. At last, substitute $X = x^2\partial$ and put $\beta_0 = 1$ to obtain, for even k , the coefficients from (3.3). If k is odd, one obtains $\beta_p = 0$ for all p .

Proposition 1 is proven. $\blacksquare$

The general form (3.2) is a consequence of Proposition 1 and decomposition (2.4).

4.2 Polynomials $P_k^n(\lambda)$. To compute the polynomials P_k^n , put $X_1 = \dots = X_n = x\partial$. One readily gets, from (3.2),

$$\sigma_\lambda(T_\lambda(X_1, \dots, X_n))|_{x=1} = n! \sum_{\substack{0 \leq k \leq n \\ k \text{ even}}} P_k^n(\lambda) \xi^{n-k}. \quad (4.1)$$

Furthermore, using the well-known expression $(x\partial)^n = \sum_{l=0}^n \left\{ \begin{smallmatrix} n \\ l \end{smallmatrix} \right\} x^l \partial^l$, one has:

$$\begin{aligned} T_\lambda(X_1, \dots, X_n) &= n! (x\partial + \lambda)^n \\ &= n! \sum_{p=0}^n \binom{n}{p} (x\partial)^n \lambda^{n-p} = n! \sum_{p=0}^n \sum_{l=0}^n \binom{n}{p} \left\{ \begin{smallmatrix} n \\ l \end{smallmatrix} \right\} x^l \partial^l \lambda^{n-p}. \end{aligned}$$

A straightforward computation gives the projectively equivariant symbol (2.5) of this differential operator:

$$\begin{aligned} \sigma_\lambda(T_\lambda(X_1, \dots, X_n))|_{x=1} &= n! \sum_{\substack{0 \leq k \leq n \\ k \text{ even}}} \sum_{p=0}^n \sum_{l=n-k}^n (l-n+k)! \frac{\binom{l}{n-k}^2 \binom{2\lambda-n+k}{l-n+k}}{\binom{n-k+l+1}{2n-2k+1}} \binom{n}{p} \left\{ \begin{smallmatrix} p \\ l \end{smallmatrix} \right\} \lambda^{n-p} \xi^{n-k}. \end{aligned}$$

Compare with the equality (4.1) to obtain the formulae from (3.4).

Theorem 1 (ii) is proven.

4.3 Uniqueness. Let T be an $\mathfrak{sl}_2$ -equivariant linear map $U(\mathfrak{sl}_2) \rightarrow \mathcal{D}_\lambda$ for a certain $\lambda \in \mathbb{C}$. In view of the decomposition (2.4), it follows from Proposition 1 that $\sigma_\lambda \circ T|_{\mathcal{F}_k} = c_k(\lambda)\mathcal{A}_k$, where $c_k(\lambda)$ is a constant depending on λ . Recall that $\text{Pol}_n(T^*\mathbb{R})$ is a *rigid* $\mathfrak{sl}_2$ -module, i.e., every $\mathfrak{sl}_2$ -equivariant linear map on $\text{Pol}_n(T^*\mathbb{R})$ is proportional to the identity (see, e.g., [15]). Assuming, now, that T preserves the principal symbol, the rigidity of $\text{Pol}_n(T^*\mathbb{R})$ fixes the constants $c_k(\lambda)$ in a unique way. Hence the uniqueness of T_λ .

Theorem 1 is proven.

5 The embedding $\mathfrak{gl}(\lambda) \rightarrow \mathcal{D}_\lambda$

A corollary of the uniqueness of the operator T_λ and results of [1, 2, 7, 17] is that the embedding $\mathfrak{gl}(\lambda) \rightarrow \mathcal{D}_\lambda$ constructed in [7] coincides with T_λ .

More precisely, according to results of [1, 2, 17], there exists a homomorphism of Lie algebras $p_\lambda : U(\mathfrak{sl}_2) \rightarrow \mathcal{D}_\lambda$ preserving the principal symbol. The homomorphism p_λ is, in particular, $\mathfrak{sl}_2$ -equivariant. By uniqueness of T_λ , one has $T_\lambda = p_\lambda$. It is also proven that the kernel of p_λ is a two-sided ideal of $U(\mathfrak{sl}_2)$ generated by $\Delta - \lambda(\lambda - 1)$ (see [1, 2]). Taking the quotient, one then has an embedding $\tilde{T}_\lambda : \mathfrak{gl}(\lambda) \rightarrow \mathcal{D}_\lambda$. Since the embedding from [7] preserves the principal symbol, it is equal to $\tilde{T}_\lambda$. Finally, it is obvious that the image of T_λ is the subalgebra $\mathcal{D}_\lambda^{\text{pol}} \subset \mathcal{D}_\lambda$ of differential operators with polynomial coefficients. Therefore, $\tilde{T}_\lambda : \mathfrak{gl}(\lambda) \rightarrow \mathcal{D}_\lambda^{\text{pol}}$ is a Lie algebras isomorphism.

6 Examples

As an illustration of Theorem 1, let us give the expressions of the general formulae (3.1) and (3.2) for the order $n = 1, 2, 3, 4, 5$. Let X_1, X_2, X_3, X_4 and X_5 be arbitrary vector fields in $\mathfrak{sl}_2$.

1) The $\mathfrak{sl}_2$ -equivariant symbol, defined by (2.5), of a first order operator of a Lie derivative $L_{X_1}^\lambda$ is

$$\sigma_\lambda(L_{X_1}^\lambda) = X_1(x)\xi.$$

2) The “anti-commutator” $[L_{X_1}^\lambda L_{X_2}^\lambda]_+$ has the following projectively equivariant symbol:

$$\sigma_\lambda([L_{X_1}^\lambda L_{X_2}^\lambda]_+) = (X_1 X_2)_+ \xi^2 + \frac{1}{3} \lambda(\lambda - 1) ((X_1' X_2')_+ - 2(X_1'' X_2)_+) \xi$$

which also following from (2.5).

3) The projectively equivariant symbol of a third order expression $[L_{X_1}^\lambda L_{X_2}^\lambda L_{X_3}^\lambda]_+$ can be also easily calculated from (2.5). The result is:

$$\begin{aligned} \sigma_\lambda([L_{X_1}^\lambda L_{X_2}^\lambda L_{X_3}^\lambda]_+) &= (X_1 X_2 X_3)_+ \xi^3 \\ &+ \frac{1}{5} (3\lambda^2 - 3\lambda - 1) ((X_1' X_2' X_3)_+ - 2(X_1'' X_2 X_3)_+) \xi. \end{aligned}$$

4) Direct calculation from (2.5) gives the projectively equivariant symbol of a fourth order expression $[L_{X_1}^\lambda L_{X_2}^\lambda L_{X_3}^\lambda L_{X_4}^\lambda]_+$, that is:

$$\begin{aligned} \sigma_\lambda([L_{X_1}^\lambda L_{X_2}^\lambda L_{X_3}^\lambda L_{X_4}^\lambda]_+) &= (X_1 X_2 X_3 X_4)_+ \xi^4 \\ &+ \frac{1}{7}(6\lambda^2 - 6\lambda - 5)((X'_1 X'_2 X_3 X_4)_+ - 2(X''_1 X_2 X_3 X_4)_+) \xi^2 \\ &+ \frac{1}{15}\lambda(\lambda - 1)(3\lambda^2 - 3\lambda - 1)((X'_1 X'_2 X'_3 X'_4)_+ - 4(X''_1 X'_2 X'_3 X_4)_+ \\ &+ 4(X''_1 X''_2 X_3 X_4)_+). \end{aligned}$$

5) In the same manner, one can easily check that the $\mathfrak{sl}_2$ -equivariant symbol of a fifth order expression $[L_{X_1}^\lambda L_{X_2}^\lambda L_{X_3}^\lambda L_{X_4}^\lambda L_{X_5}^\lambda]_+$ is:

$$\begin{aligned} \sigma_\lambda([L_{X_1}^\lambda L_{X_2}^\lambda L_{X_3}^\lambda L_{X_4}^\lambda L_{X_5}^\lambda]_+) &= (X_1 X_2 X_3 X_4 X_5)_+ \xi^5 \\ &+ \frac{5}{9}(2\lambda^2 - 2\lambda - 3)((X'_1 X'_2 X_3 X_4 X_5)_+ - 2(X''_1 X_2 X_3 X_4 X_5)_+) \xi^3 \\ &+ \frac{1}{7}(3\lambda^4 - 6\lambda^3 + 3\lambda + 1)((X'_1 X'_2 X'_3 X'_4 X_5)_+ - 4(X''_1 X'_2 X'_3 X_4 X_5)_+ \\ &+ 4(X''_1 X''_2 X_3 X_4 X_5)_+) \xi. \end{aligned}$$

Acknowledgments

I would like to thank V Ovsienko for statement of the problem. I am also grateful to Ch Duval and A El Gradechi for enlightening discussions.

References

- [1] Beilinson A and Bernstein J, Localisation de g -modules, *C.R. Acad. Sci. Paris Ser. I Math.* **292** (1981), 15–18.
- [2] Beilinson A and Bernstein J, A Proof of Jantzen Conjectures, *Adv. in Sov. Math.* **16** (1993), 1–50.
- [3] Cartan E, Leçons sur la théorie des espaces à connexion projective, Gauthier – Villars, Paris, 1937.
- [4] Cohen P, Manin Yu and Zagier D, Automorphic Pseudodifferential Operators, in *Progr. Non-linear Diff. Eq. Appl.*, Vol. 26, Birkhäuser, Boston, 1997, 17–47.
- [5] Duval C and Ovsienko V, Space of Second Order Linear Differential Operators as a Module Over the Lie Algebra of Vector Fields, *Adv. in Math.* **132**, Nr. 2 (1997), 316–333.
- [6] Dixmier J, Algèbres enveloppantes, Gauthier – Villars, Paris, 1974.
- [7] Feigin B L, The Lie Algebras $\mathfrak{gl}(\lambda)$ and Cohomologies of Lie Algebra of Differential Operators, *Russian Math. Surveys*, **43**, Nr. 2 (1988), 157–158.
- [8] Gargoubi H, Sur la géométrie de l'espace des opérateurs différentiels linéaires sur $\mathbf{R}$, *Bull. Soc. Roy. Sci. Liège* **69**, Nr. 1 (2000), 21–47.

-
- [9] Gargoubi H and Ovsienko V, Space of Linear Differential Operators on the Real Line as a Module Over the Lie Algebra of Vector Fields, *Internat. Mathem. Res. Notices* Nr. 5 (1996), 235–251.
 - [10] Gargoubi H and Ovsienko V, Modules of Differential Operators on the Real Line, *Funct. Anal. Appl.* **35**, Nr. 1 (2001), 16–22.
 - [11] Graham R, Knuth D and Patashnik O, Concrete Mathematics, Addison-Wesley, 1989.
 - [12] Grozman P and Leites D A, Lie Superalgebras of Supermatrices of Complex Size. Their Generalizations and Related Integrable Systems, in Proc. Internatnl. Symp. Complex Analysis and Related Topics, Editors: E. Ramirez de Arellano, et. al., Mexico, 1996, Birkhäuser Verlag, 1999, 73–105.
 - [13] Khesin B and Malikov F, Universal Drinfeld–Sokolov Reduction and the Lie Algebras of Matrices of Complex Size, *Comm. Math. Phys.* **175**, Nr. 1 (1996), 113–134.
 - [14] Lecomte P B A, Mathonet P and Tousset E, Comparison of Some Modules of the Lie Algebra of Vector Fields, *Indag. Math., N.S.* **7**, Nr. 4 (1996), 461–471.
 - [15] Lecomte P B A and Ovsienko V, Projectively Invariant Symbol Calculus, *Lett. Math. Phys.* **49**, Nr. 3 (1999), 173–196.
 - [16] Leites D A and Sergeev A N, Orthogonal Polynomials of a Discrete Variable and Lie Algebras of Complex-Size Matrices, *Theor. Math. Phys.* **123**, Nr. 2 (2000), 582–608.
 - [17] Shoikhet B, Certain Topics on the Representation Theory of the Lie Algebra $\mathfrak{gl}(\lambda)$. Complex Analysis and Representation Theory. 1, *J. Math. Sci. (New York)* **92**, Nr. 2 (1998), 3764–3806.
 - [18] Wilczynski E J, Projective Differential Geometry of Curves and Ruled Surfaces, Leipzig – Teubner, 1906.